


Process for preparing bisphospholane ligands

The present invention is directed at a process for the preparation of bidentate ligands based on bisphospholanes. In particular, the invention relates to the preparation of enantiomerically enriched compounds of the general formula (I) :



Enantiomerically enriched ligands are used in asymmetric synthesis or asymmetric catalysis. The important thing here is that the electronic and stereochemical properties of the ligand are optimally matched to the respective catalysis problem. An important aspect of the success of this class of compounds is believed to be the creation of an asymmetric environment around the metal centre due to these ligand systems. To utilize such an environment for effective transfer of the chirality, it is advantageous to control the flexibility of the ligand system as inherent limitation of the asymmetric induction.

Within the class of phosphorus-containing ligands, cyclic phosphines, in particular phospholanes, have attained particular importance. Bidentate, chiral phospholanes are, for example, the DuPhos and BPE ligands used in asymmetric catalysis. In the ideal case, these provide a chiral ligand framework which can be modified in a variety of ways and can be varied within a broad range in terms of its steric and electronic properties.

DE10309356 describes concrete ligand systems and a route for preparing them. The synthetic route presented there starts out from phenylphosphine which is converted into a dimetal phenolphosphine by deprotonation with a strong base
5 before the reaction to form the phospholane is carried out.


EP528865 describes the preparation of phospholanes starting from dilithiophenylphosphine and a bifunctional alkylation reagent. The preparation of the dilithiophenylphosphine is not mentioned.

10 EP1028967, JOC 2003, 68, 1701-1707 and Org. Lett. 2003, 5, 1273-75, present preparations of enantiomerically enriched phospholanes. Here too, the conversion into the phospholane is brought about using phenolphosphine or a lithiated bistrimethylsilylphosphide.

15 It was an object of the present invention to provide a further process for preparing the abovementioned phospholanes and enantiomerically enriched ligands. In particular, the process should be economical on an industrial scale from both economic and ecological points
20 of view. Very particular value should be attached to a process which starts out from materials which are readily available commercially and uses reagents which can be handled relatively unproblematically.

Such a process is proposed in the claims. Claim 1 describes
25 a process for preparing the desired ligand systems. The dependent subordinate Claims 2 to 13 are directed at preferred embodiments of the process of the invention.

Carrying out a process for preparing enantiomerically enriched compounds of the general formula (I),



where

* indicates a stereogenic centre,

R^1 and R^4 are each, independently of one another

(C_1-C_8)-alkyl, HO-(C_1-C_8)-alkyl, (C_1-C_8)-alkoxy,

5 (C_2-C_8)-alkoxyalkyl, (C_6-C_{18})-aryl, (C_7-C_{19})-aralkyl,

(C_1-C_8)-alkyl-(C_6-C_{18})-aryl, (C_3-C_8)-cycloalkyl,

(C_1-C_8)-alkyl-(C_3-C_8)-cycloalkyl,

(C_3-C_8)-cycloalkyl-(C_1-C_8)-alkyl,

R^2 and R^3 are each, independently of one another, H,

10 (C_1-C_8)-alkyl, HO-(C_1-C_8)-alkyl, (C_1-C_8)-alkoxy,


(C_2-C_8)-alkoxyalkyl, (C_6-C_{18})-aryl, (C_7-C_{19})-aralkyl,

(C_1-C_8)-alkyl-(C_6-C_{18})-aryl, (C_3-C_8)-cycloalkyl,

(C_1-C_8)-alkyl-(C_3-C_8)-cycloalkyl,

(C_3-C_8)-cycloalkyl-(C_1-C_8)-alkyl,

15 A is a C_2 bridge in which two carbon atoms have sp^2 hybridization, starting from compounds of the general formula (II),



(II)

where

R^1 to R^4 can be as defined above,

20 M is an alkali metal or a trimethylsilyl group, and reacting these with compounds of the general formula (III),




25 where

A is as defined above and

the radicals X are each, independently of one another, a nucleofugic leaving group, and preparing the compounds of the general formula (II) by reacting compounds of the


30 general formula (IV),



where

R^1 to R^4 are as defined above and

5 the radicals Y are each, independently of one another, a nucleofugic leaving group, with compounds of the general formula (V),



10

where

M is an alkali metal and Aryl is a (C_6-C_{18}) -aryl or $((C_1-C_8)$ -alkyl)₁₋₃- (C_6-C_{18}) -aryl radical, and subsequently with an alkali metal and additionally, if appropriate, with
 15 trimethylsilyl chloride, with the compounds of the formula (V) being obtained by reaction of compounds of the general formula (VI),



20


where

Aryl is as defined above,

with an alkali metal, enables the ligand systems in question to be obtained simply and particularly

25 advantageously according to the invention. It was particularly suprising that the procedure described achieves a relatively good increase in yield, which was not to have been expected from the prior art.

The process described above is preferably applied to compounds in which A is a radical from the group consisting of




5

where


- R is H, (C₁-C₈)-alkyl, (C₆-C₁₈)-aryl, (C₇-C₁₉)-aralkyl, (C₁-C₈)-alkyl-(C₆-C₁₈)-aryl, (C₃-C₈)-cycloalkyl, (C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl, and Q is O, NH, NR. Q in these formulae is particularly preferably oxygen or NR, with R being able to be (C₁-C₈)-alkyl, (C₆-C₁₈)-aryl, benzyl. Especially preferred radicals R are methyl, ethyl, propyl, isopropyl, tert-butyl, phenyl, naphthyl, fluorenyl, benzyl.
- 15 Preference is likewise given to using compounds of the formula (IV) in which R² and R³ are each H and R¹ and R⁴ are each, independently of one another, (C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl, (C₂-C₈)-alkoxyalkyl.

- 20 Preference is also given to using compounds of the general formula (III) or (IV) in which X or Y is selected from the group consisting of halogen, Otos (p-toluenesulfonate), OMes (methylsulfonate), triflate (trifluoroacetate), p-nitrobenzenesulfonate (nosylate, ONs) in the process of the invention.

Particular preference is likewise given to using compounds of the general formula (VII) or (VIII), for compounds of general formula (IV)



(VII)



(VIII)

5

where

the radicals Y are selected independently from the group consisting of halogen, OTos, OMes, triflate, nosylate, R¹ and R⁴ are each, independently of one another,

10 (C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl, (C₂-C₈)-alkoxyalkyl, (C₆-C₁₈)-aryl, (C₇-C₁₉)-aralkyl, (C₁-C₈)-alkyl-(C₆-C₁₈)-aryl, (C₃-C₈)-cycloalkyl, (C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl,

the radicals R' are each, independently of one another,

15 H, (C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl, (C₆-C₁₈)-aryl, (C₇-C₁₉)-aralkyl, (C₁-C₈)-alkyl-(C₆-C₁₈)-aryl, (C₃-C₈)-cycloalkyl, (C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl, (C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl.

Very particular preference is given to compounds of the formula (VII) or (VIII) in which R' is H, methyl, ethyl, propyl, isopropyl, tert-butyl, phenyl, and R¹ and R⁴ are each methyl, ethyl, propyl, isopropyl, tert-butyl, phenyl.

20 In principle, all elements of main group 1 of the Periodic Table can be employed as alkali metals. Preference is here given to lithium, sodium and potassium, with lithium being especially preferred as the metal to be used.

The solvents to be used for the individual steps of the process can be selected by a person skilled in the art on the basis of his general technical knowledge. The solvents should be solvents which do not if at all possible, promote

30

any secondary reactions and are themselves inert under the reaction conditions. The reaction of compounds of the general formula (VI) with alkali metals is preferably carried out in an aprotic polar solvent. Very particular
5 preference is here given to ethers such as THF, diethyl ether, dioxane, DME or DEE.

The temperature window within which the reaction according to the invention is carried out can likewise be selected freely by a person skilled in the art. Here, a person
10 skilled in the art will be guided by efficiency factors such as space-time yield, energy costs and by-products spectra and set a temperature which helps to ensure an optimal reaction. The reaction of the compound (IV) with the compound (V) is preferably carried out at a temperature
15 of from -50°C to $+100^{\circ}\text{C}$, more preferably from -30°C to $+80^{\circ}\text{C}$ and particularly preferably from -25°C to $+40^{\circ}\text{C}$. The reaction of compounds of the general formula (VI) with alkali metals can, on the other hand, be carried out at temperatures of from -25°C to $+40^{\circ}\text{C}$, preferably from -15°C
20 to $+30^{\circ}\text{C}$ and particularly preferably from -10°C to $+10^{\circ}\text{C}$.


In a further, preferred embodiment of the present process, the reactions of (VI) with alkali metals to form (V) and subsequently with (IV) and also the further reaction of the products obtained to form (III) and finally with (II) to
25 form (I) can be carried out in a single vessel. Thus, the entire reaction can be carried out in a simple fashion as a one-pot synthesis.

The present process thus offers a further very decisive advantage over the synthetic routes disclosed in the prior
30 art. Furthermore, the choice of starting substances means that no strong, difficult-to-handle bases, e.g. alkyllithium compounds, have to be used in the synthesis. As a result, the present route can be carried out on an industrial scale without costly safety equipment and


precautions, which in the final analysis helps make the products cheaper and thus economically more attractive.

The present invention is generally carried out as follows:

In step 1, the compounds of the formula (IV) are prepared
5 as shown by way of example in the following scheme.




The compounds of the general formula (V) are subsequently prepared and are then reacted with the compounds of the formula (IV).



10

After this, the further synthesis after phenyl/lithium/trimethylsilyl exchange can be completed by reaction of the compounds of the formula (III) with those of the formula (II).



15

The overall yield can reach values of > 60%, preferably > 65% and very particularly preferably > 70%, starting from compounds of the formula (VI).

For the purposes of the present invention, (C₁-C₈)-alkyl is
5 methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, hexyl, heptyl or octyl, including all bonding isomers.

The term (C₁-C₈)-alkoxy refers to (C₁-C₈)-alkyl radicals
10 which are bound via an oxygen atom to the respective molecule.

The term (C₁-C₈)-alkoxyalkyl refers to (C₁-C₈)-alkyl radicals which have an oxygen atom in their chain.

The term (C₃-C₈)-cycloalkyl encompasses cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl
15 radicals, etc.

For the purposes of the present invention, a (C₆-C₁₈)-aryl radical is an aromatic radical having from 6 to 18 carbon atoms. In particular, such radicals include phenyl, naphthyl, anthryl, phenanthryl, biphenyl radicals.

20 A (C₇-C₁₉)-aralkyl radical is a (C₆-C₁₈)-aryl radical bound via a (C₁-C₈)-alkyl radical to the molecule.

Nucleofugic leaving groups are, in particular: halogen, Otosyl (OTos), Omesyl (OMes), triflate, nosylate.

Possible halogens (Hal) are chlorine, bromine and iodine.

25 For the purposes of the invention, the term enantiomerically enriched means that the proportion of one enantiomer in the mixture with its opposite enantiomer is in a range from >50% and <100%.

The structures shown encompass all possible diastereomers and the enantiomers (R, S form) coming under the respective diastereomer.

The references cited are hereby incorporated by reference
5 into the disclosure of the present invention.

Examples :

The following examples serve to illustrate the invention. They do not in any way constitute a restriction.

Example 1: (2S,5S)-2,5-Hexanediol bismesylate

5 Variant 1:

60g of (2S,5S)-2,5-hexanediol are dissolved in 400ml of dichloromethane and 177 ml of triethylamine and the solution is cooled to 0°C. The methanesulfonyl chloride is added dropwise as a solution in 50 ml of dichloromethane,
10 with the temperature being kept below 30°C. The reaction solution is stirred at room temperature for 60 minutes and subsequently hydrolyzed with 200 ml of water. The phases are separated and the aqueous phase is extracted once with dichloromethane. The organic phase is washed with saturated
15 sodium chloride solution, dried and the solvent is removed under reduced pressure.

Yield: 138.5g; 99%

Variant 2:

20 120g of (2S,5S)-2,5-hexanediol are suspended in 450 ml of toluene and 353.5 ml of triethylamine and the mixture is cooled to 5°C. A solution of 255.9 ml of methanesulfonyl chloride in 50 ml of toluene is added dropwise to this suspension, with the temperature being kept below 30°C. The
25 reaction solution is stirred at room temperature for 1 hour and subsequently hydrolyzed with 250 ml of water. The phases are separated and the aqueous phase is extracted once with toluene. The organic phase is washed with saturated sodium chloride solution and subsequently dried
30 over magnesium sulfate. The solvent is removed under reduced pressure.

Yield: 256.7g; 92%

Example 2: (2R,5R)-2,5-Dimethyl-1-phenylphospholane

Variant 1:

5 44.4 g of lithium are suspended in 400 ml of THF and the mixture is cooled to 0°C. 143.2 g of dichlorophenylphosphine (dissolved in 200 ml of THF) are slowly added dropwise to this reaction mixture, with the temperature being kept below 30°C. The orange suspension is
10 stirred for 1 hour and the THF is subsequently removed under reduced pressure. The residue is taken up in 600 ml of dimethoxyethane and refluxed for 5 hours. After cooling to room temperature, the suspension is transferred to a further reaction vessel, with the residual lithium
15 remaining in the first vessel. The suspension is cooled to -20°C and 195 g of (2S,5S)-2,5-hexanediol bismesylate (dissolved in 200 ml of toluene) are subsequently added, with the temperature being kept below 5°C. The reaction mixture is stirred at room temperature for a further
20 8 hours and the solvent is subsequently removed under reduced pressure. The residue is taken up in 500 ml of heptane, the mixture is filtered and the solid is washed twice with 300 ml each time of heptane. The heptane is removed under reduced pressure and the crude product is
25 subsequently distilled (87-89°C / 1 mbar)

Yield: 97.1 g; 71%

¹H-NMR (C₆D₆): δ = 0.70 (dd 3H), 1.11-1.30 (m, 2H), 1.20 (dd, 3H), 1.65 (m 1H), 2.00-2.45 (m, 3H), 7.50-7.30 (5H) ppm.

30 ³¹P-NMR (C₆D₆): δ = 11.1 ppm.

Variant 2: (comparison)

375 ml of methyllithium (1.6 M in ether) are added at -20°C to a solution of 32 g of phenylphosphine in 300 ml of THF. After the addition, the solution is warmed to room temperature and stirred for 1 hour. 72.5 g of (2S,5S)-2,5-hexanediol bismesylate are dissolved in 75 ml of THF and added to the reaction solution at -20°C. The reaction mixture is warmed to room temperature and stirred for 16 hours. The solvent is subsequently removed under reduced pressure and the residue is taken up in 400 ml of heptane. The reaction mixture is filtered and the residue is washed with 200 ml of heptane. The solvent is removed under reduced pressure and the product is distilled under reduced pressure (72-75°C / 0.5 mbar).

Yield: 26.5 g; 52%

Variant 3: (comparison)

113 ml of butyllithium (1.6 M in hexane) are added at -30°C to a solution of 99 g of phenylphosphine (10% strength in hexane) in 300 ml of THF. After the addition, the solution is warmed to room temperature and stirred for 1 hour. 22.5 g of (2S,5S)-2,5-hexanediol bismesylate are dissolved in 50 ml of THF and added to the reaction solution at -30°C. The reaction mixture is warmed to room temperature and stirred for 16 hours. The solvent is subsequently removed under reduced pressure and the residue is taken up in 100 ml of heptane. The reaction mixture is filtered and the residue is washed with 100 ml of heptane. The solvent is removed under reduced pressure and the product is distilled under reduced pressure (69-75°C / 0.5 mbar).

Yield: 11.65g; 67%

Example 3: (2R,5R)-2,5-Dimethyl-1-trimethylsilylphospholane

1.67 g of lithium are suspended in 60 ml of THF. 7.75 g of (2S,5S)-2,5-dimethyl-1-phenylphospholane (dissolved in 10 ml of THF) are added dropwise at 0°C to this suspension.
5 The reaction solution is stirred at room temperature for 1 hour and subsequently transferred to a further reaction vessel, with the lithium remaining in the first vessel. 11.2 ml of chlorotrimethylsilane (dissolved in 10 ml of THF) are added to this reaction mixture. After 2 hours, the
10 solvent is removed under reduced pressure and the crude product is distilled under reduced pressure (80-90°C / 30mbar).

Yield: 5.15 g; 68%

1H-NMR (CDCl₃): δ = 0.20 (d, 9H), 1.25-1.15 (m, 6H), 2.54-
15 1.20 (m, 6H) ppm.

31P-NMR (CDCl₃): δ = -54.4 ppm.

Example 4: 2,3-Bis[(R,R)-2,5-dimethylphospholan-1-yl]maleic anhydride

20 4.71g of (2R,5R)-2,5-dimethyl-1-trimethylsilylphospholane (dissolved in 5 ml of ether) are added at 0°C to a solution of 2.09 g of dichloromaleic anhydride in 20 ml of ether and the mixture is stirred at this temperature for a further 15 minutes. After a further 30 minutes at room temperature,
25 the solution is cooled to -78°C. The product crystallizes as brown crystals. The crystals are filtered off and dried under reduced pressure.

Yield: 3.56 g; 87%

1H-NMR (CDCl₃): δ = 1.06 (dd, 6H), 1.22 (dd, 6H), 2.49-1.25
30 (m, 12H), 3.32 (m, 2H) ppm.

^{31}P -NMR (CDCl_3): $\delta = -2.2$ ppm.

Example 5: {2,3-Bis[(R,R)-2,5-dimethylphospholanyl]maleic anhydride}(cyclooctadiene)rhodium(I) tetrafluoroborate

5 1.90 g of 2,3-bis[(R,R)-2,5-dimethylphospholanyl]maleic anhydride are dissolved in 15 ml of THF and the solution is cooled to -20°C . 2.40 g of $[\text{Rh}(\text{COD})_2]\text{BF}_4$ (suspended in 20 ml of THF) are slowly added to the solution. The solution is warmed to room temperature and stirred for a
10 further 90 minutes. 25 ml of ether are added to the reaction solution and the product is subsequently filtered off. The crystals are washed with 25 ml of ether and dried under reduced pressure.

Yield: 3.51 g; 97%

15 ^1H -NMR (Aceton- d_6): $\delta = 1.23$ (dd, 6H), 1.57 (dd, 6H), 2.67-1.50 (m 18H), 3.07 (m 2H), 5.15 (s, 2H), 5.85 (s, 2H) ppm.

^{31}P -NMR (Aceton- d_6): $\delta = 63.8$ (d, $J = 151\text{Hz}$) ppm.

Example 6: 2,3-Bis[(R,R)-2,5-dimethylphospholanyl]maleimide

20 A solution of 4.50 g of (2R,5R)-2,5-dimethyl-1-trimethylsilylphospholane (dissolved in 5 ml of THF) is added dropwise at 0°C to a solution of 2.70 g of N-methyl-2,3-dibromomaleimide in 10 ml of THF and the mixture is stirred for a further one hour. The solvent and all
25 volatile components are removed under reduced pressure. The product is isolated as a red-brown oil.

Yield: 3.20g (94%)


^1H -NMR (CDCl_3): δ = 1.05 (dd, 6H), 1.22 (dd, 6H), 1.78 (m, 2H), 2.05 (m, 2H), 2.26 (m, 2H), 2.42 (m, 2H), 2.98 (d, 3H), 3.32 (m, 2H) ppm.

^{31}P -NMR (CDCl_3):

5 δ = -4.9ppm.

Claims:

1. Process for preparing enantiomerically enriched compounds of the general formula (I),



where

* indicates a stereogenic centre,

R¹ and R⁴ are each, independently of one another

(C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl, (C₁-C₈)-alkoxy,

(C₂-C₈)-alkoxyalkyl, (C₆-C₁₈)-aryl, (C₇-C₁₉)-aralkyl,

(C₁-C₈)-alkyl-(C₆-C₁₈)-aryl, (C₃-C₈)-cycloalkyl,

(C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl,

(C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl,

R² and R³ are each, independently of one another, H,

(C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl, (C₁-C₈)-alkoxy,

(C₂-C₈)-alkoxyalkyl, (C₆-C₁₈)-aryl, (C₇-C₁₉)-aralkyl,


(C₁-C₈)-alkyl-(C₆-C₁₈)-aryl, (C₃-C₈)-cycloalkyl,

(C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl,

(C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl,

A is a C₂ bridge in which two carbon atoms have sp² hybridization,

by reacting compounds of the general formula (II),



where

R¹ to R⁴ can be as defined above,

M is an alkali metal or a trimethylsilyl group,

with compounds of the general formula (III),




where

A is as defined above and

5 the radicals X are each, independently of one another,
a nucleofugic leaving group,
characterized in that the compounds of the general
formula (II) are prepared by reacting compounds of the
general formula (IV),


10



where

R¹ to R⁴ are as defined above and

15 the radicals Y are each, independently of one another,
a nucleofugic leaving group,
with compounds of the general formula (V),



20

where

M is an alkali metal and Aryl is a (C₆-C₁₈)-aryl or
(C₁-C₈)-alkyl)₁₋₃-(C₆-C₁₈)-aryl radical, and
subsequently with an alkali metal and, if appropriate,
additionally with trimethylsilyl chloride,
25 with the compounds of the formula (V) being obtained
by reaction of compounds of the general formula (VI),




where

Aryl is as defined above,
with an alkali metal.

2. Process according to Claim 1,
characterized in that

A is a radical from the group consisting of



where

R is H, (C₁-C₈)-alkyl, (C₆-C₁₈)-aryl, (C₇-C₁₉)-aralkyl,
(C₁-C₈)-alkyl-(C₆-C₁₈)-aryl, (C₃-C₈)-cycloalkyl,
(C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl,
(C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl,

Q is O, NH, NR.

3. Process according to Claim 2,
characterized in that

Q is oxygen or NR, where R can be (C₁-C₈)-alkyl,
(C₆-C₁₈)-aryl, benzyl.



4. Process according to Claim 3,
characterized in that

Q is oxygen or NR, where R can be methyl, ethyl,
propyl, isopropyl, tert-butyl, phenyl, naphthyl,
fluorenyl, benzyl.

5. Process according to one or more of Claims 1 to 4,
characterized in that

compounds of the formula (IV) in which R² and R³ are
each H and R¹ and R⁴ are each, independently of one
another, (C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl,
(C₂-C₈)-alkoxyalkyl are used.

6. Process according to one or more of Claims 1 to 5,
characterized in that
compounds of the general formula (III) or (IV) in
which X or Y is selected from the group consisting of
halogen, OTos, OMes, triflate, nosylate, are used.
7. Process according to one or more of Claims 1 to 6,
characterized in that
compounds of the general formula (VII) or (VIII),



where

the radicals Y are selected independently from the
group consisting of halogen, OTos, OMes, triflate,
nosylate,

R¹ and R⁴ are each, independently of one another,
(C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl, (C₂-C₈)-alkoxyalkyl,
(C₆-C₁₈)-aryl, (C₇-C₁₉)-aralkyl,
(C₁-C₈)-alkyl-(C₆-C₁₈)-aryl, (C₃-C₈)-cycloalkyl,
(C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl,
(C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl,

the radicals R' are each, independently of one
another,

H, (C₁-C₈)-alkyl, HO-(C₁-C₈)-alkyl, (C₆-C₁₈)-aryl,
(C₇-C₁₉)-aralkyl, (C₁-C₈)-alkyl-(C₆-C₁₈)-aryl,
(C₃-C₈)-cycloalkyl, (C₁-C₈)-alkyl-(C₃-C₈)-cycloalkyl,
(C₃-C₈)-cycloalkyl-(C₁-C₈)-alkyl,

are used for compounds of general formular (IV).

8. Process according to Claim 7,
characterized in that
R' is H, methyl, ethyl, propyl, isopropyl, tert-butyl,

phenyl, and

R¹ and R⁴ are each methyl, ethyl, propyl, isopropyl, tert-butyl, phenyl.

- 5 9. Process according to one or more of the Claims 1 to 8, characterized in that the alkali metal used is lithium.
- 10 10. Process according to one or more of Claims 1 to 9, characterized in that the reaction of compounds of the general formula (VI) with alkali metals is carried out in an aprotic polar solvent.
- 15 11. Process according to one or more of Claims 1 to 10, characterized in that the reaction of the compound (IV) with the compound (V) is carried out at a temperature of from -25°C to +40°C.
- 20 12. Process according to one or more of Claims 1 to 6, characterized in that the reaction of compounds of the general formula (VI) with alkali metals is carried out at temperatures of -10°C to +10°C.
13. Process according to Claim 1, characterized in that the reaction is carried out in a one-pot variant.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/012279

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07F9/6568

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BEILSTEIN Data, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 03/084971 A (DEGUSSA) 16 October 2003 (2003-10-16) page 13 - page 15	1-13
Y	& DE 103 09 356 A (DEGUSSA) 20 November 2003 (2003-11-20) cited in the application	1-13
Y	WO 91/17998 A (DU PONT) 28 November 1991 (1991-11-28) Scheme 1	1-13
Y	& EP 0 528 865 A (DU PONT) 3 March 1993 (1993-03-03) cited in the application	1-13
A	WO 99/24444 A (CHIROTECH TECHNOLOGY LTD) 20 May 1999 (1999-05-20) page 2 -/-	1-13

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

3 February 2005

Date of mailing of the international search report

18/02/2005

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Elliott, A

INTERNATIONAL SEARCH REPORT

Intel Application No
PCT/EP2004/012279

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	& EP 1 028 967 A (CHIROTECH TECHNOLOGY LTD) 23 August 2000 (2000-08-23) cited in the application	1-13
A	PILKINGTON C J ET AL: "Expanding the Family of Phospholane-Based Ligands: 1,2-Bis(2,5-diphenylphospholano)ethane" ORGANIC LETTERS, ACS, WASHINGTON, DC, US, vol. 5, no. 8, 17 April 2003 (2003-04-17), pages 1273-1275, XP002301264 ISSN: 1523-7060 cited in the application Scheme 1	1-13
Y	HOLZ J ET AL: "Synthesis of a New Chiral Bisphospholane Ligand for the Rh(I)-Catalyzed Enantioselective Hydrogenation of Isomeric beta-Acylamido Acrylates" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY, EASTON, US, vol. 68, no. 5, 12 February 2003 (2003-02-12), pages 1701-1707, XP002244188 ISSN: 0022-3263 cited in the application Scheme 2	1-13
Y	BURK M J: "New chiral phospholanes; synthesis, characterization, and use in asymmetric hydrogenation reactions" TETRAHEDRON: ASYMMETRY, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 2, no. 7, 1991, pages 569-592, XP002146730 ISSN: 0957-4166 cited in the application Schemes I & III	1-13
Y	BLOOMFIELD P R ET AL: "Direct preparation of phenylphosphine dilithium" CHEMISTRY AND INDUSTRY, 25 April 1959 (1959-04-25), pages 541-542, XP009005175 ISSN: 0009-3068 the whole document	1-13

-/--

INTERNATIONAL SEARCH REPORT

Intel Application No
PCT/EP2004/012279

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>YAN Y-Y ET AL: "Highly Flexible Synthetic Routes to Functionalized Phospholanes from Carbohydrates"</p> <p>JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY. EASTON, US, vol. 65, 11 February 2000 (2000-02-11), pages 900-906, XP002233658 ISSN: 0022-3263 Scheme 3</p>	1-13
Y	<p>NANDI M ET AL: "Synergistic effects of hemilabile coordination and counterions in homogeneous catalysis: new turnable monophosphine ligands for hydrovinylolation reactions"</p> <p>JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 121, no. 42, 27 October 1999 (1999-10-27), pages 9899-9900, XP002316158 ISSN: 0002-7863 Scheme I</p>	1-13
Y	<p>BRAUER D J ET AL: "Novel P,N ligands derived from (R)- and (S)-1-phenylethylamine with (2R,5R)-2,5-dimethylphospholanyl groups (DuPHAMIN) for asymmetric catalysis"</p> <p>EUROPEAN JOURNAL OF INORGANIC CHEMISTRY, no. 9, May 2003 (2003-05), pages 1748-1755, XP002316159 ISSN: 1434-1948 Scheme 1, conversion of compound 4 to 5</p>	1-13
Y	<p>LI W ET AL: "Synthesis of chiral hydroxyl phospholanes from D-mannitol and their use in asymmetric catalytic reactions"</p> <p>JOURNAL OF ORGANIC CHEMISTRY, vol. 65, no. 11, 2 June 2000 (2000-06-02), pages 3489-3496, XP002316160 ISSN: 0022-3263 Scheme I</p>	1-13

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/012279

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 03084971	A	16-10-2003	BR	0308970 A	11-01-2005
			CA	2481037 A1	16-10-2003
			DE	10309356 A1	20-11-2003
			WO	03084971 A1	16-10-2003
			EP	1490379 A1	29-12-2004
			HR	20040904 A2	31-12-2004
DE 10309356	A	20-11-2003	DE	10309356 A1	20-11-2003
			BR	0308970 A	11-01-2005
			CA	2481037 A1	16-10-2003
			WO	03084971 A1	16-10-2003
			EP	1490379 A1	29-12-2004
			HR	20040904 A2	31-12-2004
WO 9117998	A	28-11-1991	US	5008457 A	16-04-1991
			AT	116988 T	15-01-1995
			AU	652756 B2	08-09-1994
			AU	7854691 A	10-12-1991
			CA	2082166 A1	18-11-1991
			DE	69106703 D1	23-02-1995
			DE	69106703 T2	11-05-1995
			DK	528865 T3	20-03-1995
			EP	0528865 A1	03-03-1993
			ES	2067230 T3	16-03-1995
			HK	66995 A	12-05-1995
			HU	68116 A2	29-05-1995
			JP	2975683 B2	10-11-1999
			JP	5507078 T	14-10-1993
			NO	924317 A	10-11-1992
			SG	9590537 A2	01-09-1995
			WO	9117998 A1	28-11-1991
			US	5177230 A	05-01-1993
			US	5206398 A	27-04-1993
			US	5322956 A	21-06-1994
EP 0528865	A	03-03-1993	US	5008457 A	16-04-1991
			AU	652756 B2	08-09-1994
			CA	2082166 A1	18-11-1991
			DE	69106703 D1	23-02-1995
			DE	69106703 T2	11-05-1995
			EP	0528865 A1	03-03-1993
			HK	66995 A	12-05-1995
			HU	68116 A2	29-05-1995
			JP	2975683 B2	10-11-1999
			JP	5507078 T	14-10-1993
			NO	924317 A	10-11-1992
			AT	116988 T	15-01-1995
			AU	7854691 A	10-12-1991
			DK	528865 T3	20-03-1995
			ES	2067230 T3	16-03-1995
			SG	9590537 A2	01-09-1995
			WO	9117998 A1	28-11-1991
			US	5177230 A	05-01-1993
			US	5206398 A	27-04-1993
			US	5322956 A	21-06-1994
WO 9924444	A	20-05-1999	AT	206128 T	15-10-2001
			AU	1041099 A	31-05-1999

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/012279

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9924444	A	CA 2307192 A1	20-05-1999
		DE 69801841 D1	31-10-2001
		DE 69801841 T2	16-05-2002
		EP 1028967 A1	23-08-2000
		ES 2162473 T3	16-12-2001
		WO 9924444 A1	20-05-1999
		JP 2001522855 T	20-11-2001
		US 6545183 B1	08-04-2003
EP 1028967	A	23-08-2000	
		AT 206128 T	15-10-2001
		AT 233780 T	15-03-2003
		AU 1041099 A	31-05-1999
		AU 1057500 A	29-05-2000
		CA 2307192 A1	20-05-1999
		CA 2347146 A1	18-05-2000
		DE 69801841 D1	31-10-2001
		DE 69801841 T2	16-05-2002
		DE 69905746 D1	10-04-2003
		DE 69905746 T2	16-12-2004
		EP 1028967 A1	23-08-2000
		EP 1127061 A1	29-08-2001
		ES 2190266 T3	16-07-2003
		WO 0027855 A1	18-05-2000
		JP 2001522855 T	20-11-2001
		JP 2002529472 T	10-09-2002
		US 6172249 B1	09-01-2001
		US 6545183 B1	08-04-2003
		ES 2162473 T3	16-12-2001
		WO 9924444 A1	20-05-1999